

L15 ANSWER 2 OF 14 MEDLINE MEDLINE  
 ACCESSION NUMBER: 2000143207 MEDLINE  
 DOCUMENT NUMBER: 20143207 PubMed ID: 10680972  
 TITLE: **Folate**-mediated drug delivery: effect of  
 alternative **conjugation** chemistry.  
 AUTHOR: Leamon C P; DePrince R B; Hendren R W  
 CORPORATE SOURCE: Department of Oligomer Development, GlaxoWellcome Research  
 Institute, Research Triangle Park, NC 27709, USA..  
 Leamon@Endocyte.com  
 SOURCE: JOURNAL OF DRUG TARGETING, (1999) 7 (3) 157-69.  
 Journal code: 9312476. ISSN: 1061-186X.  
 PUB. COUNTRY: Switzerland  
 DOCUMENT TYPE: Journal; Article; (JOURNAL ARTICLE)  
 LANGUAGE: English  
 FILE SEGMENT: Priority Journals  
 ENTRY MONTH: 200004  
 ENTRY DATE: Entered STN: 20000421  
 Last Updated on STN: 20030118  
 Entered Medline: 20000411  
 AB When utilized as a macromolecular drug targeting ligand, folic acid  
 (Pte-Glu) has traditionally been coupled to peptides, proteins and lipids  
 via one of its two carboxylate groups fortuitously located within a  
 distal  
**glutamyl** moiety. It has been assumed in the literature that the  
**gamma-glutamyl** carboxylate of Pte-Glu is the preferred  
**conjugation** site for macromolecules enduring endocytosis via the  
**folate**-binding protein receptor. However, it is also possible that  
 the steric placement of the attached macromolecule around the vitamin's  
 pteridine moiety may be the more influential parameter controlling this  
 delivery mechanism. Using solid-phase chemistries, we have synthesized  
 dipeptide derivatives of pteric acid for the purpose of identifying the  
 preferred site onto which a macromolecule can be chemically attached  
 without compromising its endocytosis potential. Thus, using fluorescent  
 and radiolabeled **conjugates**, we have determined that  
 macromolecules attached to Pte-Glu by either an **alpha-** or  
**gamma-glutamyl** linkage could associate with  
 receptor-bearing cells at virtually identical levels. We further  
 discovered that removal of the remaining un-**conjugated**  
**glutamyl** carboxylate had no inhibitory effect on cell uptake; and,  
 the cytotoxicity of related momordin toxin **conjugates** were  
 comparable among the various pterate derivatives tested. From these  
 observations we suggest that the preparation of endocytosis-competent  
 pterate-macromolecule **conjugates** is strongly influenced by the  
 steric environment around the ligand's para-aminobenzoic acid moiety, and  
 that no selective isomeric (i.e. alphaGlu versus gammaGlu)  
**conjugation** requirement necessarily exists.